

A magneto-electric quantum wheel.

Alexander Feigel*

Soreq NRC,

Yavne 81800, Israel

(Dated: December 5, 2009)

Here we show that self-propulsion in quantum vacuum may be achieved by rotating or aggregating magneto-electric nano-particles. The back-action follows from changes in momentum of electro-magnetic zero-point fluctuations, generated in magneto-electric materials. This effect may provide new tools for investigation of the quantum nature of our world. It might also serve in the future as a “quantum wheel” to correct satellite orientation in space.

PACS numbers:

Despite some initial claims of negligible or even zero momentum transfer[1, 2], recent theoretical studies concur that material objects may acquire momentum from quantum vacuum[3, 4, 5, 6, 7, 8]. This can be explained qualitatively by considering quantum vacuum as a random fluctuating electromagnetic field (so called zero fluctuations) composed of propagating modes. Each mode possesses both energy and momentum, causing Lamb splitting of spectral levels [9, 10], Casimir attraction of objects in an empty space[11, 12, 13] and other mechanical interactions from nano to astrophysical scales[14, 15, 16]. The total momentum vanishes if the counter-propagating modes cancel mutually. It occurs in all materials except magneto-electrics, which lack both space and time symmetries[17, 18, 19].

Self-propulsion requires mechanical back-action from an external medium such as ground, water, air or even a quantum liquid[20, 21]. This can be provided by wheels or propeller-like devices. Mechanical interaction with electro-magnetic radiation can also serve as a mean for moving matter on macro- and nano-scales[22, 23, 24, 25, 26]. It seems that self-propulsion in vacuum, however, can be achieved only by a rocket-like disposal of mass, at least in foreseeable future[27].

In this article we demonstrate that aggregating or rotating magneto-electric particles change the momentum of quantum vacuum and, as a consequence they acquire the resulting difference. It follows from momentum conservation: any change in momentum of zero fluctuations is compensated by a corresponding change in the momentum of a material object or electromagnetic field. These new occurrences of the vacuum momentum transfer do not require external means, such as previously proposed modification of the magneto-electric constant by applying external electric and magnetic fields[3, 8] or suppressing the quantum vacuum modes by cavity-imposed boundary conditions[4, 5].

Consider an object in crossed electric E_x and magnetic B_y fields (respectively in x and y directions). It will

experience a Lorentz force F given by:

$$F \propto B_y \frac{\partial P_x}{\partial t}, \quad (1)$$

where P is its average electric polarisation. For the sake of simplicity, P will include only the linear dielectric ϵ and magneto-electric χ terms:

$$P_x \propto \epsilon E_x + \chi_{xy} B_y, \quad (2)$$

Merging eqs. (1) and (2) results in:

$$F \propto B_y \frac{\partial \epsilon E_x}{\partial t} + \chi_{xy} \frac{1}{2} \frac{\partial B_y^2}{\partial t} + B_y^2 \frac{\partial \chi_{xy}}{\partial t}. \quad (3)$$

The first term describes a linear dielectric in external fields and includes no quantum vacuum contribution. The second and the third terms describe both classical and quantum phenomena, since $\langle B^2 \rangle$ does not vanish due to zero fluctuations of electro-magnetic field.

According to eq. (3), momentum can be extracted from quantum vacuum by four different approaches. One way is to suppress zero fluctuations by applying some dynamic boundary conditions (by a shrinking cavity for instance) which eliminate the low frequency modes of quantum vacuum, modifying in this way the $\langle B_{vac}^2 \rangle$ value[4, 5]. In this case F will be given by:

$$F_{vac} \propto \chi_{xy} \frac{1}{2} \frac{\partial \langle B_{vac}^2 \rangle}{\partial t}. \quad (4)$$

Another way is based on achieving a non-zero value for the time derivative of χ_{xy} , resulting in:

$$F_{vac} \propto \langle B_{vac}^2 \rangle \frac{\partial \chi_{xy}}{\partial t}. \quad (5)$$

The magneto-electric constant χ_{xy} consists of intrinsic and induced terms:

$$\chi_{xy} = \chi_{xy}^0 + \kappa_1 E_x B_y + \kappa_2 E_x + \kappa_3 B_y, \quad (6)$$

where χ_{xy}^0 is the intrinsic magneto-electricity, while κ_1 , κ_2 and κ_3 are the responses to external fields. This leads to a second approach by using time dependant external electric and magnetic fields[3, 8].

*Electronic address: sasha@soreq.gov.il

The third, previously unforeseen, method to extract momentum is based on rotation of the electro-magnetic material. The magneto-electric constant is a tensor and its value may be changed, therefore, by rotation. The corresponding force is:

$$F_{vac} \propto B_{vac}^2 > \frac{\partial \chi_{xy}^0}{\partial t}. \quad (7)$$

For instance, rotation by π around the x axis changes the sign of χ_{xy}^0 . In this case, the maximum momentum transfer is obtained by maximum change of the magneto-electric constant $\propto 2\chi_{xy}^0$.

The total velocity gain of a magneto-electric particle rotated by π in quantum vacuum is:

$$\Delta v_{rot} \approx A \frac{\hbar}{\rho} \frac{2\chi_{xy}^0}{a^4} \approx A \frac{\hbar}{m} \frac{2\chi_{xy}^0}{a}, \quad (8)$$

where $A \approx 10^{-2}$ is a numerical constant, ρ is the density, a is the size and $m \propto \rho a^3$ is the mass of the particle. The size defines the highest frequencies ω_{cut} of quantum vacuum contributing to the particle's momentum[3]. The modes with wavelength λ ($\lambda = c/\omega$, where c is velocity of light) which are smaller than a do not contribute to the net polarization P_x of the object and, therefore, vanish in (1). The same expression follows from a change in momentum of quantum vacuum

$$p_{vac} \approx A\hbar\chi/a, \quad (9)$$

derived by the rigorous renormalization formalism[5, 7, 8].

Another method for momentum transfer may be based on separation or aggregation of the magneto-electric particles while preserving their orientation. Momentum of quantum vacuum in N magneto-electrics of size a is greater than in a single object of size $L = N^{1/3}a$ (see (9)). The difference in momenta corresponds to the velocities:

$$\Delta v_{agg} \propto A \frac{\hbar}{\rho} \chi_{xy}^0 \left(\frac{1}{a^4} - \frac{1}{L^4} \right). \quad (10)$$

Aggregation of the particles affects the boundary conditions of the high-frequency vacuum modes generating greater forces than a shrinking external cavity, which modifies only the low-frequency ones.

An experimental investigation of momentum transfer in quantum vacuum would be preferably based on rotation or aggregation of magneto-electric materials. In these configurations the contribution of quantum vacuum can be easily isolated, while with alternating electric or magnetic fields[3, 8] the quantum and classical forces are difficult to distinguish, see (3).

The discussed effect does not interfere with other quantum effects at nano-scales. Quantum fluctuations of the position or of the magneto-electric constant of particles do not affect the average value of their momentum, as a consequence of the conservation of momentum law.

A propulsion engine may be designed by using for instance an addressable array of small magneto-electric particles or wires. Rotating (see Fig. 1) or aggregating (see Fig. 2) these particles will result in velocity:

$$\Delta V \propto \Delta v \frac{m}{M}, \quad (11)$$

where Δv is given either by (8) or by (10), m and M are the masses of the rotated particles and the entire load correspondingly. Although the proposed engine will consume energy for manipulation of the particles, the propulsion will occur without any loss of mass.

To correct the attitude of a satellite, several degrees per day are required[28]. This corresponds to a tangential velocity of $1\mu\text{m}/\text{sec}$ at 1m radius. This value may be obtained with a magneto-electric constant of $\chi_{me}^0 \approx 10^{-3}$ and $\approx 1\text{nm}$ particles, assuming that roughly half of the satellite mass is composed of magneto-electric material (see eqs. (8) and (11) taking $\rho \approx 1\text{g}/\text{cm}^3$). Materials with magneto-electric constants up to 10^{-4} at nanometric scale have been already reported, for instance FeGaO_3 in a weak ($\approx 100\text{Oe}$) external magnetic field[29]. One may reasonably assume that this gap will be closed in the future due to investigations in new composites[19] and advances in micro-patterning techniques[30].

To conclude, mechanical action of quantum vacuum on magneto-electric objects may be observable and have a significant value. Rotation or self-assembly of the nanoparticles is enough to generate a back-action from zero electro-magnetic fluctuations. The amount of momentum that can be extracted from quantum vacuum by this effect, may have in the future practical implications, depending on advances in magneto-electric materials.

-
- [1] van Tiggelen, B. A. & Rikken, G. L. J. A. Comment on "quantum vacuum contribution to the momentum of dielectric media". *Phys. Rev. Lett.* **93**, 268903 (2004).
 [2] Obukhov, Y. N. & Hehl, F. W. Forces and momenta caused by electromagnetic waves in magnetoelectric media. *Phys. Lett. A* **372**, 3946–3952 (2008).
 [3] Feigel, A. Quantum vacuum contribution to the momentum of dielectric media. *Phys. Rev. Lett.* **92**, 020404

- (2004).
 [4] van Tiggelen, B. A., Rikken, G. L. J. A. & Krstic, V. Momentum transfer from quantum vacuum to magneto-electric matter. *Phys. Rev. Lett.* **96**, 130402 (2006).
 [5] Birkeland, O. J. & Brevik, I. Feigel effect: Extraction of momentum from vacuum? *Phys. Rev. E* **76**, 066605 (2007).
 [6] Shen, J. Q. Momentum transfer between quantum vac-

- uum and anisotropic medium. *Prog. Theor. Phys.* **119**, 351–360 (2008).
- [7] van Tiggelen, B. A. Zero-point momentum in complex media. *Europ. Phys. J. D* **47**, 261–269 (2008).
- [8] Kawka, S. & van Tiggelen, B. A. Quantum electrodynamics of casimir momentum. *Arxiv* 0908.4390 (2009).
- [9] Lamb Jr., W. E. & Retherford, R. Fine structure of the hydrogen atom by a microwave method. *Phys. Rev.* **72**, 241–243 (1947).
- [10] Bethe, H. A. The electromagnetic shift of energy levels. *Phys. Rev.* **72**, 339–341 (1947).
- [11] Casimir, H. B. G. On the attraction between two perfectly conducting plates. *Proc. Kon. Ned. Akad. Wetensch.* **51**, 793795 (1948).
- [12] Lamoreaux, S. K. Demonstration of the casimir force in the 0.6 to 6 μm range. *Phys. Rev. Lett.* **78**, 5–8 (1997).
- [13] Bressi, G., Carugno, G., Onofrio, R. & Ruoso, G. Measurement of the casimir force between parallel metallic surfaces. *Phys. Rev. Lett.* **88**, 041804 (2002).
- [14] Davies, P. C. W. Quantum vacuum friction. *J. Opt. B* **7**, S40–S46 (2005).
- [15] Capasso, F., Munday, J. N., Iannuzzi, D. & Chan, H. B. Casimir forces and quantum electrodynamical torques: Physics and nanomechanics. *IEEE J. Sel. Top. Quantum Electron.* **13**, 400–414 (2007).
- [16] Dupays, A., Rizzo, C., Bakalov, D. & Bignami, G. F. Quantum vacuum friction in highly magnetized neutron stars. *Epl* **82**, 69002 (2008).
- [17] Fiebig, M. Revival of the magnetoelectric effect. *J. of Phys. D* **38**, R123–R152 (2005).
- [18] Eerenstein, W., Mathur, N. D. & Scott, J. F. Multiferroic and magnetoelectric materials. *Nature* **442**, 759–765 (2006).
- [19] Arima, T. Magneto-electric optics in non-centrosymmetric ferromagnets. *J. of Phys. Cond. Mat.* **20**, 434211 (2008).
- [20] Dreyfus, R. *et al.* Microscopic artificial swimmers. *Nature* **437**, 862–865 (2005).
- [21] Avron, J. E., Gutkin, B. & Oaknin, D. H. Adiabatic swimming in an ideal quantum gas. *Phys. Rev. Lett.* **96**, 130602 (2006).
- [22] Arcizet, O., Cohadon, P. F., Briant, T., Pinard, M. & Heidmann, A. Radiation-pressure cooling and optomechanical instability of a micromirror. *Nature* **444**, 71–74 (2006).
- [23] Eichenfield, M., Camacho, R., Chan, J., Vahala, K. J. & Painter, O. A picogram- and nanometre-scale photonic-crystal optomechanical cavity. *Nature* **459**, 550–U79 (2009).
- [24] Groblacher, S., Hammerer, K., Vanner, M. R. & Aspelmeyer, M. Observation of strong coupling between a micromechanical resonator and an optical cavity field. *Nature* **460**, 724–727 (2009).
- [25] Hugel, T. *et al.* Single-molecule optomechanical cycle. *Science* **296**, 1103–1106 (2002).
- [26] Kippenberg, T. J. & Vahala, K. J. Cavity optomechanics: Back-action at the mesoscale. *Science* **321**, 1172–1176 (2008).
- [27] Millis, M. G. & Davis, E. W. (eds.) *Frontiers of Propulsion Science (Progress in Astronautics and Aeronautics)* (AIAA, 2009).
- [28] Montenbruck, O. & Gill, E. *Satellite Orbits: Models, Methods and Applications* (Springer, 2005).
- [29] Kubota, M. *et al.* X-ray directional dichroism of a polar ferrimagnet. *Phys. Rev. Lett.* **92**, 137401 (2004).
- [30] Kida, N. *et al.* Enhanced optical magnetoelectric effect in a patterned polar ferrimagnet. *Phys. Rev. Lett.* **96**, 167202 (2006).
-




FIG. 1: **Propulsion in quantum vacuum by rotation of magneto-electric particles.** Rotation of magneto-electric particles will modify the momentum of quantum vacuum, and generate a corresponding back-action. The resulting velocity is proportional to intrinsic value of the magneto-electric tensor χ_{me}^0 .




FIG. 2: **Propulsion in quantum vacuum by aggregation of magneto-electric particles.** Changes in dimensions of an object with intrinsic χ_{me}^0 will affect the momentum of quantum vacuum, generating a back-action. The corresponding velocity increases strongly with smaller particle sizes.